


Pneumo Update Europe 2018

15 - 16 June, Budapest

**Hot Topic: Pulmonary Imaging-
What the scientific world is
talking about**



Hans-Ulrich Kauczor, Germany

What is new? What is important?


- **Technical Advances**
- **Airway diseases: COPD, CF**
- **Diffuse parenchymal lung disease**
- **Pulmonary nodules, lung cancer, radiomics**
- **Precision cancer therapy: pulmonary complications**

Technical advances

State-of-the Art: Technical Advances

- **Plain film, projection radiography**
- **CT: helical, dual energy, spectral**
- **MRI:**
 - proton: ultrashort TE, contrast agent: Gd, oxygen**
 - non-proton: hyperpolarized gases: ^3He , ^{129}Xe**


Digital Tomosynthesis




- 158 pulmonary nodules
- tomosynthesis: 120kVp, 60 images, 30° rotation, copper filter, dose 0.1 mSv
- x-ray: two planes; 120kVp; dose 0.06 mSv



X-ray




Tomosynthesis




CT

Dobbins-JT et al. Radiology 2017; 282:236-250
Ferrari-A et al. Ann Transl Med 2018;6:91

Digital Tomosynthesis




➤ 158 pulmonary nodules






Dobbins et al. Radiology 2017; 282:236-250
Ferrari-A et al. Ann Transl Med 2018;6:91

Digital Tomosynthesis as good as CT?

1111 participants of
Swedish cardiopulmonary
Bioimage Study (SCAPIS)
125 with 252 nodules




*Meltzer-C et al. Radiology 2018;
epub*




Darkfield Radiography

➤ X-Rays, but scatter instead of attenuation; porcine model




Photon Counting CT

- Lower noise, higher resolution



- Improved further by high resolution matrix




Symons-R et al. Radiology 2017;285:980-989
Pourmorteza-A et al Invest Radiol 2018; 53:365-372

μ -CT: Fibrosis


- Fibrosis starts peripherally and grows centrally in the lobule
- Alveolar collapse might be trigger of fibroproliferation




clinical



μ postmortem



clinical




μ postmortem

Mai-C et al. Radiology 2017;283:252-263

μ -CT: Fibrosis







- Separation of aerated volume, tissue volume, conducting airways
- TNF transgenic mice have higher tissue volume



*Bell-RD et al. PLoS One
2018;13:e0190678*

μ -CT: Fibrosis

- Separation of aerated volume, tissue volume, conducting airways
- TNF transgenic mice have higher tissue volume



*Bell-RD et al. PLoS One
2018;13:e0190678*

Amid-proton transfer MRI

- 82 patients with pulmonary nodules, NH- groups of mobile proteins and peptides; comparison DWI; PET/CT





TABLE 1. Differences in Differentiation Between Malignant and Benign Nodules for Each Index

	Malignant nodules (Mean \pm SD)	Benign nodules (Mean \pm SD)	<i>P</i> value
Age	70.8 \pm 6.8	69.7 \pm 6.5	0.19
Long axis diameter*	18.5 \pm 5.4	16.9 \pm 5.9	0.29
MTR _{asym} (3.5 ppm) (%)	2.0 \pm 6.4	-2.9 \pm 4.9	0.0002
ADC ($\times 10^{-3}$ mm ² /s)	1.2 \pm 0.3	1.3 \pm 0.2	0.0024
SUV	2.2 \pm 1.6	2.3 \pm 0.5	0.0008

Phase-resolved ventilation and perfusion MRI


- Regional mapping, fourier decomposition
- Feasibility in volunteers COPD, CTEPH and CF patients



Voskrebenzev-A et al. *Magn Reson Med*
2018;79:2306-2314

Phase-resolved ventilation and perfusion MRI

- Regional mapping, fourier decomposition
- Feasibility in volunteers COPD, CTEPH and CF patients



Take Home: Technical Advances

- **More innovations to be expected**
- **Radiography => tomosynthesis**
- **Radiography => dark field**
- **CT => photon counting, dark field ?**
- **MRI => CEST, non-contrast, non-proton
=> functional, molecular**

Airway diseases

COPD, CF


State-of-the Art: Airway disease

- **CT => high resolution, low dose**
- **MRI => high resolution, low dose**
- **Quantification, imaging biomarkers**
- **Remodeling, exacerbation**
- **Surveillance**

COPD: Visual Emphysema


- 3171 COPDGene participants
inspiratory CT
obs agreement 0.7-0.85
correlation with
- increasing airflow
 - dyspnea
 - symptoms
 - GOLD stage
 - mortality

Lynch-DA et al. Radiology 2018 in press





COPD: CT Biomechanics

- 490 COPDGene participants
in-/expiratory CT
registration
deformation field
better prediction of
- resp. quality of life
 - BODE index
 - mortality






COPD: CT vs MRI

- 224 lung cancer screening participants
- Low dose CT
- Propeller T2W MRI



Meier-Schroers-M et al. Fortschr Röntgenstr 2018; epub




COPD: Imaging Biomarkers



Kirby-M et al. *Eur J Radiol* 2017;86: 335-342


COPD: Ventilation MRI

- 27 patients
- ^{19}F MRI
- Free breathing
- Dynamic
- Defect volume
- Wash-out time
- Fractional ventilation




*Gutberlet-M et al.
Radiology
2018;296:1040-1051*

Cystic Fibrosis: Airway Remodeling (CT)



**Konietzke.O et al. PLoS
2018; April-9
Weinheimer-O et al
Proc SPIE Med Imag
2017**


Cystic Fibrosis: Airway Remodeling (CT)



*Konietzke.O et al. PLoS
2018; April-9
Weinheimer-O et al
Proc SPIE Med Imag
2017*

Cystic Fibrosis: Exacerbation (MRI)




- 56 patients; 29 with exacerbations
- Discrepancies: T2 = structure, DWI = function (cellularity)



Ciet-P et al. Eur Resp J 2017;50:1601437

Cystic Fibrosis: Surveillance (MRI)






- 97 patients; MRI vs. LCI: complementary outcome measures
- MRI: regional perfusion and ventilation inhomogeneity



Stahl-M et al. *Am J Respir Crit Care Med* 2017;195:349-359

Cystic Fibrosis: Non-contrast MRI vs. CT



➤ 29 patients, in/ex CT and proton MRI



Pennati-F et al. J Magn Reson Imaging 2018; epub

Cystic Fibrosis: Non-contrast vs. CE-MRI


- 50 patients, T2w + contrast-enhanced T1w MRI
- Mosaic perf score



Leutz-Schmidt-P et al. *Eur J Radiol*
2018; 101:178-183

Cystic Fibrosis: Oxygen-enhanced UTE MRI



- 18 subjects, healthy, asthma, CF, repeatability



Zha-W et al. *J Magn Reson Imaging* 2018;47:1287-1297

Cystic Fibrosis: Oxygen-enhanced UTE MRI

- 18 subjects, healthy, asthma, CF, repeatability



Zha-W et al. *J Magn Reson Imaging* 2018;47:1287-1297

Take Home: Airway Diseases

- **CT => high resolution, low dose**
- **MRI => high resolution, no dose**
- **Quantification, imaging biomarkers**
- **Remodeling, exacerbation**
- **Surveillance**

Diffuse Parenchymal Lung Disease

State-of-the Art:


Diffuse Parenchymal Lung Disease

- **ATS/ERS guidelines**
- **Central role of CT**
- **MDT board diagnosis**
- **Recent white paper of the Fleischner Society**

Lynch-DA et al. Lancet Respir Med 2018;6:138-153


Interstitial Lung Abnormalities in Smokers

- Danish lung cancer screening trial
- Interstitial lung abnormalities in 332 (17%) participants
- Increased mortality regardless of fibrotic or non-fibrotic phenotype
- partly due to an association with lung cancer and non-pulmonary malignancies




Interstitial Lung Abnormalities in Diabetics

- 48 non-diabetics
68 patients with pre-diabetes
29 newly diagnosed
110 patients with long-term T2D
- metabolic control
diabetes-related complications
breathlessness
lung function
- T2D: increased breathlessness,
higher prevalence of restrictive
lung disease,
CT evidence of interstitial disease
- => diabetic pneumopathy




Idiopathic Pulmonary Fibrosis: CT

- Pattern analysis and vascular volume
- “Severity”: more GGO, more reticulation => poor outcome
- Vessel-related structure predicts survival better FVC decline



Idiopathic Pulmonary Fibrosis: CT

- 144 IPF patients
- Pattern analysis
- Fibrosis score independent predictor of survival => poor outcome




Lee-SM et al. *Eur Radiol* 2018;28:1293-1300


Idiopathic Pulmonary Fibrosis: PET/CT

- 113 IPF patients; FDG-PET/CT
- Target-to-background SUV < 4.9 med. survival 71 mo
> 4.9 med. survival 24 mo


Win-T et al. Eur J Nuc Med Mol Imag 2018;45:806-815



a




b





Pulmonary Fibrosis: Gadofosveset-enhanced MRI

- Vascular leakage
- Gadofosveset is a FDA-approved albumin-binding contrast agent
- Detects vascular permeability, used for vascular MRI
- 6 pulmonary fibrosis patients; FDG-PET/CT
- Vascular leak increased vs. health controls




Pulmonary Fibrosis: Iron-oxid-enhanced MRI

- Superparamagnetic iron-oxide nanoparticles (ferumoxytol)
- Iron uptake in macrophages
- UTE MRI (CODE)
- Enhancement in granulomatosis disease, not cancer





Yoon-SH et al. KRJ 2018;19:153-157



Pulmonary Fibrosis: Hypox 129Xe Gas Exchange MRI






➤ 13 healthy, 12 IPF patients



Wang-JM et al. Thorax 2018;73:21-28

Pulmonary Fibrosis: Hypx 129Xe Gas Exchange MRI

➤ 13 healthy, 12 IPF patients



Wang-JM et al. Thorax 2018;73:21-28

Take Home: Diffuse Parenchymal Lung Disease

- **Comorbidities**
- **Quantification, imaging biomarkers**
- **Activity, remodeling:** metabolism
vascular permeability
inflammation (macrophage)

Pulmonary Nodules

Lung Cancer

Radiomics

State-of-the-Art: Pulmonary Nodules, Lung Cancer

➤ Fleischner Guidelines 2.0

A: Solid Nodules*

Nodule Type	Size		
	<6 mm (<100 mm ³)	6–8 mm (100–250 mm ³)	>8 mm (>250 mm ³)
Single			
Low risk [†]	No routine follow-up	CT at 6–12 months, then consider CT at 18–24 months	Consider CT at 3 months, PET/CT, or tissue sampling
High risk [†]	Optional CT at 12 months	CT at 6–12 months, then CT at 18–24 months	Consider CT at 3 months, PET/CT, or tissue sampling


MacMahon-H et al. Radiology 2017;284:228-243

➤ Lung Rads

➤ Lung cancer staging 8th edition



State-of-the Art: Radiomics

- Quantitative imaging analytics
- Precision diagnostics, predictive modeling of outcome



Pulmonary Nodules: Fleischner Guidelines 2.0



- 2066 patients with coronary CTA, 578 pulmonary nodules
- Patients required follow-up reduced by 65%



Scholtz-JE et al. *J Cardiovasc Comput Tomogr* 2018;12:28-33


Pulmonary Nodules: Radiomics Prediction Model

- 72 pulmonary nodules (31 benign, 41 malignant)
- Lung Image Database Consortium image collection
- Accuracy 85%, better than Lung Rads 72%



Pulmonary Nodules: Radiomics Prediction Model


- 72 pulmonary nodules (33 benign, 42 malignant)
- Retrospective analysis of clinical routine CT
- Accuracy 84%, sensitivity 93%, specificity 73%



Chen-CH et al. PLoS One 2018 Feb-5

Lung Adenocarcinoma Heterogeneity: Radiomics



- 763 patients: radiomics + genomics
- EGFR, KRAS mutations



Rios Velazquez-E et al. *Cancer Res* 2017;77:3922-3930

Pulmonary Nodules: MRI

- 51 patients
- 119 pulmonary nodules
- CT and UTE MRI



Part-solid	20 (23.6)	17 (13.5)	15 (37.7)	82 (37.0)
Pure ground-glass opacity	11 (10.0)	12 (10.9)	10 (90.9)	97 (98.0)

Wielpütz-MO et al. AJR 2018;210:1-10
Ohno-Y et al. JMRI 2018 epub




TABLE 4. Currently Recommended MR Protocols for Nodule Detection and Evaluation

		MR technique	Additional comments
Nodule detection		STIR turbo SE or HASTE imaging	Black-blood technique would be better to be applied. Respiratory gating would be better to be applied rather than breath holding, if possible.
		Pulmonary thin-section MRI with ultra-short TE	Ultra-short TE would be better to be shorten at $\leq 192 \mu s$. Respiratory gating would be better to be applied rather than breath holding, if possible.
Nodule evaluation	Nondynamic technique	T1-weighted turbo SE or HASTE imaging	Black-blood technique would be better to be applied. Respiratory gating would be better to be applied rather than breath holding, if possible.
		3D T1-weighted radiofrequency-spoiled GRE imaging with interpolation and/or partial Fourier techniques as well as fat suppression method	VIBE (Siemens Healthcare), LAVA (GE Healthcare), THRIVE (Philips Medical Systems), Quick 3D (Canon < previously Toshiba > Medical Systems)
		T2-weighted turbo SE or HASTE imaging	Black-blood technique would be better to be applied. Respiratory gating would be better to be applied rather than breath holding, if possible.
		STIR turbo SE or HASTE imaging	Black-blood technique would be better to be applied. Respiratory gating would be better to be applied rather than breath holding, if possible. Moreover, contrast ratio between lesion to muscle would be better to be evaluated for quantitative assessment.
		Diffusion-weighted imaging	Apparent diffusion coefficient as well as lesion-to-spinal cord ratio evaluations would be better to be applied.
		Contrast-enhanced T1-weighted turbo SE or HASTE imaging	Black-blood technique would be better to be applied. Respiratory gating would be better to be applied rather than breath holding, if possible.
		Contrast-enhanced 3D T1-weighted radiofrequency-spoiled GRE imaging with interpolation and/or partial Fourier techniques as well as fat suppression method	VIBE (Siemens Healthcare), LAVA (GE Healthcare), THRIVE (Philips Medical Systems), Quick 3D (Canon < previously Toshiba > Medical Systems)
	Dynamic contrast-enhanced technique	Dynamic first-pass contrast-enhanced T1-weighted GRE imaging with ultra-fast TE	Ultra-short TE would be better to be shorten at ≤ 0.2 msec. Temporal resolution would be better to be set at ≤ 1.5 sec with and without applying parallel imaging technique. Bolus injection protocol would be better to be considered with field strength.

VIBE: volumetric interpolated breath-hold examination, LAVA: liver acquisition with volume acceleration, THRIVE: T1 high resolution isotropic volume excitation, Quick 3D: fast and segmented 3D T1-weighted spoiled-gradient-echo.


Lung Cancer Heterogeneity: DWI Cell Count

- ## ➤ Feasibility, imaging histology pipeline, validation



Lung Cancer Staging: PET/MRI vs. PET/CT

- 77 patients: comparable staging results, therapeutic decisions



Schaarschmidt-BM et al. Eur Radiol 2017;27:681-688


Take Home: Nodules, Lung Cancer, Radiomics

- **Tumor characterization**
- **Tumor heterogeneity**
- **Therapy response** **early detection**
 quantification
 prediction

Precision Cancer Therapy: Pulmonary Complications

State-of-the-Art: Pulmonary Complications


- Immunotherapy
- Checkpoint inhibitors: PD-1, PD-L1, CTLA-4



- Immune-related adverse events (irAEs): pneumonitis, hypophysitis, thyroiditis, hepatitis, pancreatitis, colitis

Pulmonary Complications of Immunotherapy


➤ Toxic pneumonitis, different patterns



Rituximab-CHOP AIP/DAD pattern



**Nivolumab
NSIP pattern**




Nivolumab AIP/DAD pattern




**Nivolumab
HP pattern**

Pulmonary Complications of Immunotherapy


- Toxic pneumonitis, COP pattern



Nivolumab



Nivo off, steroids on



Steroids off

Pulmonary Complications of Immunotherapy

- Toxic pulmonary edema
- Toxic pleural effusion



Nilotinib




Nivolumab + Ipilimumab

Pulmonary Complications of Immunotherapy


➤ Sarcoid mimikry

Lymphadenopathy



Ipilimumab

Perilymphatic nodules



Ipilimumab

NSCLC Treatment: PD1-Inhibitors ir RECIST

- Immune-related RECIST > RECIST 1.1, e.g. longer TTP
RR = 18 %; pseudoprogression rare





Fig. 2 Time to progression by irRECIST and RECIST1.1



Nishino-M et al. J Immunotherapy Cancer 2016;4:84;
Nishino-M et al. Clin Cancer Res 2017;23:5737-5744

NSCLC Treatment: PD1-Inhibitors ir RECIST

- Immune-related RECIST > RECIST 1.1, e.g. longer TTP
RR = 18 %; pseudoprogression rare






Fig. 2 Time to progression by irRECIST and RECIST1.1



Nishino-M et al. J Immunotherapy Cancer 2016;4:84;
Nishino-M et al. Clin Cancer Res 2017;23:5737-5744

Take Home: Pulmonary Complications



List of References

1. Dobbins JT, Mcadams HP, Sabol JM, Chakraborty DP, Kazerooni EA, Reddy GP, Vikgren J, Bath M. Multi-institutional evaluation of digital tomosynthesis, dual energy radiography and conventional chest radiography for the detection and management of pulmonary nodules. *Radiology* 2017; 282:236-250
2. Ferrari A, Bertolaccini L, Solli P, Di Salvia PO, Scaradozzi D. Digital chest tomosynthesis: the 2017 updated review of an emerging application. *Ann Transl Med* 2018;6:91
3. Meltzer C, Vikgren J, Bergman B, Molnar D, Norrlund RR, Hassoun A, Gottfridsson B, Båth M, Johnsson AA. Detection and characterization of solid pulmonary nodules at digital chest tomosynthesis: data from a cohort of the pilot Swedish Cardiopulmonary Bioimage Study. *Radiology* 2018 epub
4. Gromann LB, De Marco F, Willer K, Noël PB, Scherer K, Renger B, Gleich B, Achterhold K, Fingerle AA, Muenzel D, Auweter S, Hellbach K, Reiser M, Baehr A, Dmochewitz M, Schroeter TJ, Koch FJ, Meyer P, Kunka D, Mohr J, Yaroshenko A, Maack H, Pralow T, van der Heijden H, Proksa R, Koehler T, Wieberneit N, Rindt K, Rummeny EJ, Pfeiffer F, Herzen J. In-vivo X-ray Dark-Field Chest Radiography of a Pig. *Sci Rep* 2017;7:4807

List of References

5. Hellbach K, Baehr A, De Marco F, Willer K, Gromann LB, Herzen J, Dmochewitz M, Auweter S, Fingerle AA, Noël PB, Rummeny EJ, Yaroshenko A, Maack HI, Pralow T, van der Heijden H, Wieberneit N, Proksa R, Koehler T, Rindt K, Schroeter TJ, Mohr J, Bamberg F, Ertl-Wagner B, Pfeiffer F, Reiser MF. Depiction of pneumothoraces in a large animal model using x-ray dark-field radiography. Sci Rep 2018;8:2602
6. Hellbach-K, Meinel FG, Conlon TM, Willer K, Yaroshenko A, Velroyen A, Braunagel M, Auweter S, Reiser MF, Eickelberg O, Pfeiffer F, Yildirim AÖ. X-Ray Dark-field Imaging to Depict Acute Lung Inflammation in Mice. Sci Rep. 2018 Feb 1;8(1):2096.
7. Symons R, Pourmorteza A, Sandfort V, Ahlman MA, Cropper T, Mallek M, Kappler S, Ulzheimer S, Mahesh M, Jones EC, Malayeri AA, Folio LR, Bluemke DA Feasibility of Dose-reduced Chest CT with Photon-counting Detectors: Initial Results in Humans Radiology. 2017 Dec;285(3):980-989
8. Pourmorteza A, Symons R, Henning A, Ulzheimer S, Bluemke DA Dose Efficiency of Quarter-Millimeter Photon-Counting Computed Tomography: First-in-Human Results Invest Radiol. 2018 Jun;53(6):365-372.

List of References

9. Mai C, Verleden SE, McDonough JE, Willems S de Wever W, Coolen J Dubbeldam A, van Raemdonck DE, Verbeken EK, Verleden GM Hogg JC, Vanaudenaerde BM, Wuyts WAS, Verschakelen JA. Thin-section CT features of idiopathic pulmonary fibrosis correlated with micro-CT and histologic analysis. *Radiology* 2017;283:252-263
10. Bell RD, Rudmann C, Wood RW, Schwarz EM, Rahimi H. Longitudinal micro-CT as an outcome measure of interstitial lung disease in TNF-transgenic mice. *PLoS One* 2018;13:e0190678
11. Ohno Y, Kishida Y, Seki S, Yui M, Miyazaki M, Koyama H, Yoshikawa T. Amide proton transfer-weighted imaging to differentiate malignant from benign pulmonary lesions: comparison with diffusion-weighted imaging and FDG-PET/CT. *JMRI* 2018;47:1013-1021
12. Voskresenzov A, Gutberlet M, Klimeš F, Kaireit TF, Schönfeld C, Rotärmel A, Wacker F, Vogel-Claussen J. Feasibility of quantitative regional ventilation and perfusion mapping with phase-resolved functional lung (PREFUL) MRI in healthy volunteers and COPD, CTEPH, and CF patients. *Magn Reson Med*. 2018 Apr;79(4):2306-2314.

List of References

13. Lynch DA, Moore CM, Wilson CG, Nevrekar D, Jennermann T, Humphries SM, Austin JHM, Grenier PA, Kauczor HU, Han MLK, Regan EA, Make BJ, Russell P, Bowler RP, Beaty TH, Curran-Everett D, Hokanson JE, Curtis JL, Silverman EK, Crapo JD. CT-based Visual Classification of Emphysema: Association with Mortality in the COPDGene Study. Radiology 2018 in press
14. Bodduluri S, Bhatt SP, Hoffman ES, Newell JD, Martinez CH, Dransfield MT, Han MK, Reinhardt JM. Biomechanical CT metrics associated with patient outcomes in COPD. Thorax 2017;72:409-414
15. Meier-Schroers M, Sprinkart AM, Becker M, Homsy R, Thomas D. Quantitative and Qualitative Assessment of Pulmonary Emphysema with T2-Weighted PROPELLER MRI in a High-Risk Population Compared to Low-Dose CT. Fortschr Rontgenstr 2018 epub
16. Kirby M, van Beek EJR, Seo JB, Biederer J, Nakano Y, Coxson HO, Parraga G. Management of COPD: is there a role for quantitative imaging? Eur J Radiol 2017;86: 335-342
17. Gutberlet M, Kaireit TF, Voskrebenezv A, Lasch F, Freise J, Welte T, Wacker F, Hohlfeld JM, Vogel-Claussen J. Free-breathing Dynamic ¹⁹F Gas MR Imaging for Mapping of Regional Lung Ventilation in Patients with COPD. Radiology 2018 Mar;286(3):1040-1051.

List of References

18. Konietzke P, Weinheimer O, Wielpütz MO, Savage D, Ziyeh T, Tu C, Newman B, Galbán CJ, Mall MA, Kauczor HU, Robinson TE. Validation of automated lobe segmentation on paired inspiratory-expiratory chest CT in 8-14 year-old children with cystic fibrosis. PLoS One. 2018;13(4):e0194557
19. Weinheimer O, Wielpütz M, Konietzke P, Heussel C, Kauczor HU, Brochhausen C, Hollemann D, Savage D, Galban C, Robinson T. Fully automated lobe-based airway taper index calculation in a low dose MDCT CF study over 4 time-potins. Proc SPIE Med Imag 2017;10133
20. Ciet P, Bertolo S, Ros M, Andrinopouzlou E, Tavano V, Lucca F, Feiweier T, Krestin GP, Tiddens HAWM, Morana G. Detection and monitoring of lung inflammation in cystic fibrosis during respiratory tract exacerbation using diffusion-weighted MRI. Eur Resp J 2017;50:1601437
21. Stahl M, Wielpütz MO, Graeber SY, Joachim C, Sommerburg O, Kauczor HU, Puderbach M, Eichinger M, Mall MA: Comparison of lung clearance index and MRI for assessment of lung disease in children with cystic fibrosis. AJRCCM 2017;195:349-359
22. Pennati F, Roach DJ, Clancy JP, Brody AS, Fleck RJ, Aliverti A, Woods JC. Assessment of pulmonary structure-function relationships in young children and adolescents with cystic fibrosis by multivolume proton-MRI and CT. J Magn Reson Imaging. 2018 Feb 19.

List of References

23. Leutz-Schmidt P, Stahl M, Sommerburg O, Eichinger M, Puderbach MU, Schenk JP, Alrajab A, Triphan SMF, Kauczor HU, Mall MA, Wielpütz MO. Non-contrast enhanced magnetic resonance imaging detects mosaic signal intensity in early cystic fibrosis lung disease. *Eur J Radiol.* 2018 Apr;101:178-183.
24. Zha W, Kruger SJ, Johnson KM, Cadman RV, Bell LC, Liu F, Hahn AD, Evans MD, Nagle SK, Fain SB. Pulmonary ventilation imaging in asthma and cystic fibrosis using oxygen-enhanced 3D radial ultrashort echo time. *J Magn Reson Imaging.* 2018 May;47(5):1287-1297.
25. Lynch DA, Sverzellati N, Travis WD, Brown KK, Colby TV, Galvin JR, Goldin JG, Hansell DM, Inoue Y, Johkoh T, Nicholson AG, Knight SL, Raoof S, Richeldi L, Ryerson CJ, Ryu JH, Wells AU. Diagnostic criteria for idiopathic pulmonary fibrosis: a Fleischner Society White Paper. *Lancet Respir Med.* 2018 Feb;6(2):138-153.
26. Hoyer N, Wille MMW, Thomsen LH, Wilcke T, Dirksen A, Pedersen JH, Saghir Z, Ashraf H, Shaker SB. Interstitial lung abnormalities are associated with increased mortality in smokers. *Respir Med.* 2018 Mar;136:77-82.
27. Kopf S, Jan B, Groener JB, Zoltan Kender Z, Fleming T, Brune M, Riedinger C, Volk N, Herpel E, Pesta D, Szendroedi J, Wielpütz MO, Kauczor HU, Katus H, Kreuter M, Nawroth PP. Breathlessness and restrictive lung disease: an important diabetes-related feature in patients with type 2 diabetes. *Respiration* 2018 in press

List of References

28. Jacob-J et al. Bartholmai BJ, Egashira R, Brun AL, Rajagopalan S, Karwoski R, Kokosi M, Hansell DM, Wells AU. Chronic hypersensitivity pneumonitis: identification of key prognostic determinants using automated CT BMC Pulm Med 2017;17:81
29. Jacob-J Bartholmai BJ, Rajagopalan S, Karwoski R, Mak SM, Mok W, Della Casa G, Sugino K, Walsh SLF, Wells AU, Hansell DM. Automated computer-based CT stratification as a predictor of outcome in hypersensitivity pneumon et al. Eur Radiol 2017; 27:3635-3646
30. Jacob J, Bartholmai BJ, Rajagopalan S, van Moorsel CHM, van Es HW, van Beek FT, Struik MHL, Kokosi M, Egashira R, Brun AL, Nair A, Walsh SLF, Cross G, Barnett J, de Lauretis A, Judge EP, Desai S, Karwoski R, Ourselin S, Renzoni E, Maher TM, Altmann A, Wells AU. Predicting Outcomes in Idiopathic Pulmonary Fibrosis Using Automated CT analysis. Am J Respir Crit Care Med. 2018 Apr 23 epub
31. Lee SM, Seo JB, Oh SY, Kim TH, Song JW, Lee SM, Kim N. Prediction of survival by texture-based automated quantitative assessment of regional disease patterns on CT in idiopathic pulmonary fibrosis. Eur Radiol. 2018 Mar;28(3):1293-1300.

List of References

32. Win T, Screaton NJ, Porter JC, Ganeshan B, Maher TM, Fraioli F, Endozo R, Shortman RI, Hurrell L, Holman BF, Thielemans K, Rashidnasab A, Hutton BF, Lukey PT, Flynn A, Ell PJ, Groves AM. Pulmonary 18F-FDG uptake helps refine current risk stratification in idiopathic pulmonary fibrosis. *Eur J Nucl Med Mol Imaging*. 2018 May;45(5):806-815.
33. Montesi SB, Rao R, Liang LL, Goulart HE, Sharma A, Digumarthy SR, Shea BS, Seethamraju RT, Caravan P, Tager AM. Gadofosveset-enhanced Lung MRI to Detect Ongoing Vascular Leak in Pulmonary fibrosis. *Eur Respir J*. 2018 Apr 5 epub
34. Yoon SH, Lee C, Park J, Goo JM, Park JY. In-vivo Visualization of Iron Oxide Enhancement in Focal Pulmonary Inflammatory Lesions Using a Three-Dimensional Radial Gradient-Echo-Based Ultrashort Echo Time Sequence: A Preliminary study. *Korean J Radiol*. 2018;19(1):153-157.
35. Wang JM, Robertson SH, Wang Z, He M, Virgincar RS, Schrank GM, Smigla RM, O'Riordan TG, Sundy J, Ebner L, Rackley CR, McAdams P, Driehuys B. Using hyperpolarized 129Xe MRI to quantify regional gas transfer in idiopathic pulmonary fibrosis. *Thorax*. 2018 Jan;73(1):21-28.

List of References

36. MacMahon H, Naidich DP, Goo JM, Lee KS, Leung ANC, Mayo JR, Mehta AC, Ohno Y, Powell CA, Prokop M, Rubin GD, Schaefer-Prokop CM, Travis WD, Van Schil PE, Bankier AA. Guidelines for Management of Incidental Pulmonary Nodules Detected on CT Images: From the Fleischner Society 2017. *Radiology*. 2017 Jul;284(1):228-243.
37. Davatzikos C, Rathore S, Bakas S, Pati S, Bergman M, Kalarot R, Sridharan P, Gastounioti A, Jahani N, Cohen E, Akbari H, Tunc B, Doshi J, Parker D, Hsieh M, Sotiras A, Li H, Ou Y, Doot RK, Bilello M, Fan Y, Shinohara RT, Yushkevich P, Verma R, Kontos D. Cancer imaging phenomics toolkit: quantitative imaging analytics for precision diagnostics and predictive modeling of clinical outcome. *J Med Imaging (Bellingham)*. 2018 Jan;5(1):011018.
38. Scholtz JE, Lu MT, Hedgire S, Meyersohn NM, Oliveira GR, Prabhakar AM, Gupta R, Kalra MK, Shepard JO, Hoffmann U, Ghoshhajra BB. Incidental pulmonary nodules in emergent coronary CT angiography for suspected acute coronary syndrome: Impact of revised 2017 Fleischner Society Guidelines. *J Cardiovasc Comput Tomogr*. 2018;12(1):28-33.
39. Choi W, Oh JH, Riyahi S, Liu CJ, Jiang F, Chen W, White C, Rimner A, Mechalakos JG, Deasy JO, Lu W. Radiomics analysis of pulmonary nodules in low-dose CT for early detection of lung cancer. *Med Phys*. 2018 Apr;45(4):1537-1549

List of References

40. Chen CH, Chang CK, Tu CY, Liao WC, Wu BR, Chou KT, Chiou YR, Yang SN, Zhang G, Huang TC. Radiomic features analysis in computed tomography images of lung nodule classification. PLoS One. 2018 Feb 5;13(2)
41. Rios Velazquez-E, Parmar C, Liu Y, Coroller TP; Cruz G, Stringfield O, Ye Z, Makrigiorgios M, Fenessy F, Mak RH, Gillies R, Quakenbush J, Aerts HJWL. Somatic mutations drive distinct imaging phenotypes in lung cancer. Cancer Res 2017;77:3922-3930
42. Wielpütz MO, Lee HY, Koyama H, Yoshikawa T, Seki S, Kishida Y, Sakai Y, Kauczor HU, Sugimura K, Ohno Y. Morphological Characterization of Pulmonary Nodules With Ultrashort TE MRI at 3 T. AJR 2018; Mar 16:1-10
43. Ohno Y, Kauczor HU, Hatabu H, Seo JB, van Beek EJR; International Workshop for Pulmonary Functional Imaging (IWPF). MRI for solitary pulmonary nodule and mass assessment: Current state of the art. J Magn Reson Imaging. 2018 Mar 23.
44. Yin Y, Sedlacek O, Muller B, Warth A, Gonzalez-Vallinas M, Lahrmann B, Grabe N, Kauczor HU, Breuhahn K, Vignon-Clementel IE, Drasdo D. Tumor Cell Load and Heterogeneity Estimation From Diffusion-Weighted MRI Calibrated With Histological Data: an Example From Lung Cancer. IEEE Trans Med Imaging. 2018 Jan;37(1):35-46.

List of References

45. Schaarschmidt BM, Grueneisen J, Metzenmacher M, Gomez B, Gauler T, Roesel C, Heusch P, Ruhlmann V, Umutlu L, Antoch G, Buchbender C. Thoracic staging with 18F-FDG PET/MR in non-small cell lung cancer - does it change therapeutic decisions in comparison to 18F-FDG PET/CT? *Eur Radiol*. 2017 Feb;27(2):681-688.
46. Nishino M, Hatabu H, Sholl LM, Ramaiya NH. Thoracic Complications of Precision Cancer Therapies: A Practical Guide for Radiologists in the New Era of Cancer Care. *Radiographics*. 2017 Sep-Oct;37(5):1371-1387.
47. Nishino M, Ramaiya NH, Chambers ES, Adeni AE, Hatabu H, Jänne PA, Hodi FS, Awad MM. Immune-related response assessment during PD-1 inhibitor therapy in advanced non-small-cell lung cancer patients. *J Immunother Cancer*. 2016 Dec 20;4:84
48. Nishino M, Dahlberg SE, Adeni AE, Lydon CA, Hatabu H, Jänne PA, Hodi FS, Awad MM. Tumor Response Dynamics of Advanced Non-small Cell Lung Cancer Patients Treated with PD-1 Inhibitors: Imaging Markers for Treatment Outcome. *Clin Cancer Res*. 2017 Oct 1;23(19):5737-5744